

Marine Birds – Scoter Species Pre-migration Staging Surveys

description

This atlas page illustrates the best available data that identifies Surf, Black, White-winged and Unidentified Scoters pre-migration staging areas, which falls within the months of March through to May. The Surf Scoter (*Melanitta perspicillata*) is a large, stocky duck with white patches on its face. It breeds on lakes in boreal forests and tundra, and is commonly seen on the coast of British Columbia during winter months. The Black Scoter (*Melanitta nigra*) is darker in appearance, breeds on lakes in the sub-arctic, and migrates to coastal areas during winter months. The White-winged scoter (*Melanitta fusca*) has white patches on its wings, breeds furthest inland compared to the other Scoter species, and is most often seen on inland rivers and lakes during migration. All three scoter species feed primarily on aquatic invertebrates, particularly molluscs.

Source data used to develop this map consisted of a range of shoreline-based, ground, boat and aerial surveys, which were focused on identifying, counting and recording marine birds. The majority of the surveys were collected at sites with defined boundaries and areas. Surveys collected along transects from each dataset were buffered depending upon observation width described within the source survey methodology in order to generate a survey area.

Observations of the listed species were totalled for each survey location and divided by the area surveyed to generate a density value for each survey location. The densities of all of the survey locations within a single planning unit were averaged and adjusted for survey effort to calculate the illustrated metric, average density of Surf, Black, White-winged, and Unidentified Scoters in pre-migration staging by planning unit. The values were then classified and symbolized using 5 quantiles, meaning that each class consisted of 20% of the non-zero planning units. (A quantile is established by dividing the frequency distribution of a variable into equal groups: that is, each quantile contains the same fraction of the total number of values being measured.)



PHOTO: RICH ENGELBRECHT

data sources

- Environment Canada (Canadian Wildlife Service) - BC Ferry Surveys
- Environment Canada (Canadian Wildlife Service) – Burrard Inlet Environmental Action Plan: Winter Bird Surveys
- Environment Canada (Canadian Wildlife Service) – Coastal Waterbird Inventory
- Environment Canada (Canadian Wildlife Service) – West Coast Vancouver Island Waterbird Survey
- Bird Studies Canada - BC Coastal Waterbird Survey
- Alan Burger - Southwest Vancouver Island Shelf Surveys
- Capital Regional District – Harbours Atlas
- Laskeek Bay Conservation Society - Laskeek Bay Surveys

data resolution

- Weighted averages were calculated for each 2 kilometre x 2 kilometre planning unit.

data collected

- 1939-2008

date compiled

- 2010

reviewers

- Peter Davidson, Bird Studies Canada
- James Kenyon, Ducks Unlimited Canada
- Representatives from Environment Canada, Canadian Wildlife Service

reviewer comments

- It might have been appropriate to separate out the scoters into the separate species.

caveats of use

- Survey effort is not consistent across all planning units, across all areas of the coast or across all seasons. Areas with no data may not have been surveyed and these data gaps are not necessarily indicative of an absence of Scoter species.
- The density values in an area can change over time in response to natural population fluctuations and changes in habitat conditions (natural or anthropogenic).
- The precision of the average density values displayed in the legend is not indicative of the original precision of the survey counts.
- This feature is a compilation of data collected by many people, for different purposes, using different survey techniques with different methodologies within each technique and, therefore, considerable care must be taken when using the data.
- Surveys performed in different seasons will produce different results.
- Recommended date of expiry for use of these data in a marine planning context: None provided.

map, feature data and metadata access

- Visit www.bcmca.ca/data for more information.

references

Species description contains material from:

- B.C. Conservation Data Centre. *Species Summary: Melanitta perspicillata, Melanitta nigra and Melanitta fusca*. BC Ministry of Environment (MoE) 2010. 21 Sep. 2010 <http://a100.gov.bc.ca/pub/eswp/>
- The Cornell Lab of Ornithology. *All about Birds*. 2009. www.allaboutbirds.org/guide

BCMCA Atlas
Marine Birds
Scoter Species
Pre-migration Staging Surveys

Legend

Birds per square kilometre standardized by effort

- 0.01 - 1.61
- 1.62 - 5.37
- 5.38 - 13.99
- 14.00 - 40.08
- 40.09 - 11969.93

Note:
 - Classification based on 5 quantiles.

Data Sources:

Environment Canada
 (Canadian Wildlife Service),
 Bird Studies Canada,
 Alan Burger,
 Capital Regional District,
 Laskeek Bay Conservation Society



Base Data:

ESRI Base Data, GeoBase, GeoBC,
 NOAA, Natural Resources Canada,
 USGS, Washington State Government

Thematic Data:

For more information on data sources
 and methods please refer to the
 facing page to this map

Projection: BC Albers NAD83



1:4,250,000 *



* Written scales are approximate and are based on a 11 x 17 inch paper size.

Prepared for:



Map template by Caslys Consulting Ltd.

August 27, 2010



Note: The inset has been provided to illustrate the level of detail available in the data. The inset is for illustrative purposes only and does not imply any significance to this region.